



Zinx Lozenges
zincum aceticum 2x

Zinx GCP(guafenesin/carbetapentane citrate/phenylephrine hydrochloride)
Oral Solution
100 mg/15 mg/5 mg per 5 mL

Rx Only

Zinx Lozenges are used to reduce the number of days your cold lasts and to treat the symptoms associated with the common cold. Zinx is a lozenge that is designed to dissolve slowly in your mouth. **DO NOT CRUSH OR CHEW.**

Zinx GCP is used to treat symptoms of cough and congestion associated with the common cold, asthma, bronchitis, and other upper respiratory tract conditions. You should seek treatment from your physician if you have a primary disease or condition such as asthma or an upper respiratory tract infection that is causing the coughing and congestion. Zinx GCP is solution that is taken by mouth.

- **You should not take/give Zinx Lozenges or Zinx GCP if . . .**

Your are under 6 years of age, are a nursing mother, are taking or have taken any medication known as a monoamine oxidase inhibitor (MAOI) in the last 14 days, or you have sensitivity or are allergic to any of the ingredients. If you are not sure if the medication you are taking an MAOI, please ask your doctor.

- **Warning**

Do not eat or drink fruit or fruit juices for 30 minutes before or after taking a Zinx Lozenge because this can cause the lozenge to not work as well.

Use Zinx GCP with caution if you have high blood pressure, heart disease, a thyroid condition, diabetes, glaucoma, or have an enlarged prostate.

Do not drink alcohol, including beer, wine, and liquor, or perform dangerous activities while taking Zinx GCP due to the possibility of feeling drowsy as a result of taking the medicine.

Avoid contact with your eyes. Flush your eyes with water if you accidentally get either Zinx or Zinx GCP in your eyes.

- **Side Effects**

Some patients taking Zinx Lozenges have noticed a bad aftertaste, chalkiness, or irritation inside the mouth. Some patients taking the ingredients in Zinx GCP have experienced drowsiness, dry mouth, nervousness, excitability, headache and upset stomach.

This listing of side effects is not complete. Your healthcare professional can discuss with you a complete list of potential side effects that can occur when the products in the Zinx Cough Kit.

- **What I Should Know About Taking Zinx Cough Kit with Other Medication . . .**

Zinx Lozenges are not is not known to interfere with the other medications.

If you are taking or have taken an MAOI within the last 14 days, the MAOI may cause a greater and longer response to Zinx GCP than intended.

Always tell your healthcare provider if you are taking the components of Zinx Cough Kit or other medications so that he/she is fully aware. Always talk to your healthcare provider before starting any new medications.

ZCHKCPI-01

Revised: Mar 07

Auriga Pharmaceuticals, LLC

An Auriga Laboratories Company

Norcross, GA 30092